



Prevalence and Correlates of Medication Non-adherence among Hypertensive Patients on Follow-up in Central Kenya

Ernest M. Mutua¹, Okello B.M Agina², Isaac J. Mwanzo³

1. Dedan Kimathi University of Technology, School of Health Sciences, Dedan Kimathi, Nyeri, Kenya
2. Kenyatta University, School of Medicine, Nairobi, Kenya
3. Kenyatta University, School of Public Health & Applied Human Sciences, Nairobi, Kenya

Abstract

Background and Objective: In Kenya, a significant number of hypertensive patients have uncontrolled blood pressure, but data on medication adherence among this patient population is scarce. This study sought to determine the prevalence and correlates of medication non-adherence among hypertensive patients on follow-up in two referral hospitals in central Kenya.

Methods: We undertook a cross-sectional study comprising of 339 hypertensive patients undergoing follow-up care in two referral hospitals located in Nyeri County between October and December 2019. Medication adherence was assessed using a validated medication adherence questionnaire. Medication adherence was a dichotomous variable (adherent vs non-adherent), with the prevalence of non-adherence being measured as a proportion. Independent predictors of medication non-adherence were identified by fitting a multiple logistic regression model, where adjusted odds ratios (AORs) were computed for various covariates and interpreted at a 5% level of significance and 95% confidence interval (CI).

Results: The study sample largely comprised of the female gender (65.2%) and elderly people (mean age, 65 years +/- 12). Nearly half (46.6%) of the hypertensive patients interviewed were non-adherent to their medications. The factors associated with non-adherence were: having poor hypertension knowledge (AOR, 5.6, 95% CI, 3.3, 9.4); being on more than one antihypertensive medication (AOR, 2.8, 95% CI, 1.7, 4.7) and being on two or more daily doses of medications (AOR, 2.3, 95% CI, 1.3, 4.1).

Conclusion and Implications for Translation: Non-adherence to prescribed medications is highly prevalent among hypertensive patients in central Kenya, and it should therefore be identified as a public health concern. Strategies aimed at optimizing medication adherence among hypertensive patients in this region may consider focusing on educational interventions to increase hypertension knowledge and also on simplification of treatment regimens.

ARTICLE HISTORY

Received: 12/11/2025

Accepted: 25/11/2025

Published: 10/12/2025

KEYWORDS

Hypertension
Non-adherence
Prevalence
Correlates
Antihypertensive
Medication
Kenya

CORRESPONDENCE

Ernest M. Mutua

e-mail:

ernmutua@gmail.com

Introduction

Hypertension is the leading modifiable risk factor of cardiovascular morbidity and all-cause mortality worldwide.^{1,2} In 2019, it was estimated that over one billion people globally had hypertension, with the number representing a double increase in prevalence since 1990.³ In

Kenya, a nationally representative survey conducted in 2015 reported that 28.6% of adults aged 18-69 years were living with hypertension.⁴ Existing evidence highlights the need for achieving adequate blood control among people living with hypertension in order to minimize the risk of developing end-organ



complications such as heart disease, stroke, and kidney disease.⁵

The current Kenya Ministry of Health clinical guidelines on the management of hypertension recommend that patients with the condition should aim to achieve a blood level of <140/90 mmHg.⁶ However, achieving this clinical target has remained a challenge, with available data indicating that only 12.5% of people living with hypertension in Kenya have their blood pressures within the recommended limits.⁴ It is therefore important that barriers contributing to poor blood pressure control are identified and addressed in order to optimize treatment outcomes among people living with hypertension.

Non-adherence to medication among hypertensive patients has been identified as one of the most important barriers to blood pressure control.⁷ The World Health Organization defines adherence as the extent to which a persons' behavior (including taking of prescribed medications) corresponds with agreed recommendations from a healthcare provider.⁷ A previous systematic review comprising of studies conducted in multiple countries estimated that nearly half of patients with hypertension are non-adherent to their prescribed medications.⁸

In Kenya, Published studies on prevalence of medication non adherence among hypertensive patients are scarce, and there is no national data to quantify the burden of the problem. The few studies that have been done indicate that between 43% and 85.4% of hypertensive patients have suboptimal medication adherence⁹⁻¹² The factors contributing to non-adherence in the Kenyan context have also not been conclusively addressed in the existing studies. Synthesis of research conducted elsewhere indicates that non-adherence to antihypertensive medication is influenced by several determinants, including patient's knowledge about hypertension, affordability of medications and therapy related factors such as complexity of treatment regimens and medication adverse effects, among other factors.¹³

To our knowledge, there is no study that has been Published regarding medication adherence among hypertensive patients on follow-up in Nyeri County, Kenya. This study therefore sought to estimate the prevalence of non-adherence to antihypertensive medications and explore the associated factors in order to inform the design of interventions to address the problem in the Kenyan healthcare setting

Methods

Study Design

We conducted a cross sectional study to determine the prevalence of medication non-adherence and the associated factors among hypertensive patients on follow-up in two tertiary care hospitals in central Kenya between October and December, 2019. The study was ethically reviewed and approved by Kenyatta University Ethics Review Committee [Reference Number: KU/ERC/APPROVAL/VOL.1 (267)].

Study Setting and Participants

This study was conducted in two tertiary hospitals located in Nyeri County, Kenya: Nyeri County Referral Hospital (NCRH) and Consolata Mathari Mission Hospital (CMMH). Nyeri County is one of the 47 counties in Kenya, located in the central region of the Country. Nyeri County Referral Hospital is a level 5 hospital that offers both general and specialized medical and surgical services to Nyeri County residents. Consolata Mathari Mission Hospital is a faith-based health care organization run by the Catholic Archdiocese of Nyeri, Kenya. The hospital offers both inpatient and outpatient services.

Nyeri County is among four counties that were included in the Country's National Universal Health Coverage (UHC) pilot program launched in 2018.¹⁴ Under UHC, all patients visiting the different County health facilities were provided with services free of charge. During the survey period, hypertensive patients on follow-up in Nyeri County Referral Hospital were accessing medical services, including clinical consultation

and prescribed medications, without making any co-payment. The eligible participants for this study were hypertensive patients currently receiving any antihypertensive medication; confirmation of hypertension diagnosis and medication use was done through chart review. Patients who were newly diagnosed with hypertension at the time of the survey were excluded from the study, given that questions on medication adherence would only apply to patients who had been initiated on treatment previously.

Sample Size and Sampling

The primary objective of this study was to determine the prevalence of medication non-adherence among hypertensive patients on follow-up in the two selected tertiary care hospitals in Nyeri County. Therefore the approach for calculation of the desired minimum sample size was based on the formula for estimation of a proportion as has been described elsewhere in the literature.¹⁵

The computation of the sample size was implemented using the *open-epi* software¹⁶ incorporating the following ingredients: estimated prevalence of non-adherence among hypertensive patients of 37.6%, based on a previous hospital based study in Kenya¹²; margin of error of 5%; confidence level of 95%; and population size of N=5000 derived from the total caseload of hypertensive patients seen in the two hospitals in the preceding year before the survey of 2018. Based on these parameters, a total of 313 patients was determined as the minimum sample size. This figure was further adjusted upwards for a possible non-response rate of 10%, yielding a total of 348 patients.

The total sample size was distributed to the two hospitals based on their proportionate contribution to the combined caseload of patients seen in the preceding year before the survey. For Consolata Mathari Mission Hospital, the percentage contribution to the total caseload was 19%, thus translating to a sub-sample size of 66 patients. Patients in Nyeri County Referral Hospital accounted for 81% of the total

caseload, hence translating into a sub-sample size of 282 patients. The patients to be interviewed on a given clinic day were enrolled through systematic sampling. Patients visiting the regular clinics in each hospital are usually booked on a specific day of the week. The list of patients booked for each clinic was used as the sampling frame, with the K^{th} interval being determined by dividing the total number of booked patients by the number of patients targeted for an interview on the material day. The index case was determined through simple random sampling of the first K^{th} patients as captured in the clinic list of booked patients.

Study Variables and Data Collection

The dependent variable for this study was medication adherence which was measured using a validated medication adherence questionnaire (MAQ).¹⁷ MAQ is a self-report tool comprising of four yes/no questions: 1) Do you at times forget to take your medicines for hypertension? 2) Are there times when you feel there is no need of taking your medicines for hypertension? 3) When you feel better, do you sometimes stop taking your medicines for hypertension? and 4) Sometimes if you feel worse when you take your medicines for hypertension, do you stop taking them? A response of “Yes” in each question is awarded one point, while “No” is allocated zero. Patients answering “Yes” to at least one question (score ≥ 1) are categorized as non-adherent, while those answering “No” to all questions (score=0) are considered adherent.

Knowledge about hypertension was assessed through a set four questions developed after a review of the literature to identify the specific aspects of hypertension that are poorly understood in relation to medication adherence.^{18,19} The questions included: 1) Do you know the blood pressure reading that indicates good control of hypertension? (If yes, specify); 2) Do you know of any health problems that can arise from having high blood pressure? (If yes, specify); 3) Can someone be having high blood pressure without experiencing any symptoms? and 4) For how long are you going to be taking your medicines for hypertension?

Each correct response per question was awarded a score of one, with the total maximum score being four. The total score obtained per participant was converted into a percentage, and a cut-off of 70% was set to dichotomize the knowledge level into two categories: poor knowledge (<70%) and good knowledge ($\geq 70\%$). This categorization was adopted from a previous related study.²⁰ The data collection tools also captured information on socio-demographic characteristics and clinical details, including, diagnosis and prescribed medications. Data was collected by trained research assistants under supervision of the study investigators. Eligible participants were first taken through the details of the study, and then subsequently invited to sign an informed consent form if willing to participate. All the above survey tools- medication adherence and the knowledge assessment questionnaires were framed in two languages- English and Swahili. The original versions which were in English, were translated to Swahili language by a Kenyan language expert who was fluent in both languages. Back translation of the Swahili versions was also done and similarity checks undertaken thereafter to ensure comparability between the two versions. Patients participating in the survey were given the option of choosing their preferred language for interview.

Statistical Analysis

Statistical analysis for this study was conducted using STATA 11.2 statistical software (Stata Corp, Texas, USA). Descriptive analysis was undertaken using summary statistics including mean and standard deviation for continuous variables, and proportions for categorical variables. Analysis of factors associated with medication adherence was undertaken in two steps. Firstly, bivariate analysis was done using chi square and fisher's exact tests to explore any association between the different independent variables (socio-demographic and clinical characteristics) and medication adherence. In step two, covariates were selected for possible inclusion in multivariate analysis at a cut off of $p < 0.2$. The dependent variable- medication adherence- was dichotomized into two

outcomes: adherent and non-adherent. Multivariate analysis was undertaken by fitting a multiple logistic regression model using the stepwise forward selection method at a 5% probability of entry. The adjusted odds ratios (AORs) generated from the output were interpreted using p values and 95% confidence intervals (CI).

Results

Characteristics of Study Sample

Out of a total of 348 hypertensive patients targeted for the baseline survey, 339 of them participated, translating into a response rate of 97.4%. The participants predominantly comprised of older persons, with a mean age of 65 years (+/- 12). The female gender accounted for a large majority of the participants, comprising of about two thirds (65.2%) of the study sample.

Prevalence of Medication Non-adherence

Nearly half (46.6%) of the patients included in the survey were non-adherent to their medications based on self-report assessment. On subgroup analysis, men appeared to have a higher level of medication non-adherence, though this variation was not statistically significant.

Hypertension-Related Knowledge

Close to two thirds (59.9%) of the hypertensive patients surveyed had poor knowledge about hypertension. Knowledge about the recommended blood pressure target was strikingly low, with only 18.3% being able to correctly state the values for good control. A majority of the patients (60.8%) also lacked an understanding that blood pressure can be elevated in an individual who is not experiencing any symptoms of illness. A significant number of the participants (43.7%) were unable to name at least one hypertension mediated organ complication such as kidney disease, heart disease or stroke. The findings in this survey also indicated a lack of understanding about the lifelong nature of hypertension treatment; close to half (46.6%) of those interviewed did not know that they were

supposed to take their medications throughout unless advised to stop by their healthcare providers.

Correlates of Medication Non-adherence

On bivariate analysis, factors that were associated with medication adherence at 5% significance level, included knowledge about hypertension, number medications prescribed and number of daily doses (Table 2). At multivariate level, medication non-adherence was independently predicted by poor hypertension related knowledge, being on two or more medications and having a regimen comprising of twice or more daily dosing. Patients who had poor knowledge of hypertension were 5.6 times more likely to be non-adherent to their medications compared to those with good knowledge. The odds of being non-adherent to medications was 2.8 times higher for patients on two or more medications compared to those on mono-therapy. Also, the likelihood of being no-adherent was 2.3 times higher for patients receiving two or more doses per day compared to those on single daily doses.

Discussion

Our findings showed that nearly half (46.6%) of hypertensive patients on follow in the two referral hospitals in central Kenya had sub-optimal adherence to their prescribed medications. This finding is consistent with a previous cross sectional study that documented a 42.4% level of medication non-adherence among hypertensive patients on follow-up in a tertiary hospital in the coastal region of Kenya.²¹ However, other studies conducted previously in different settings in Kenya recorded a higher level of medication non-adherence.^{10,22} This variation may be attributed to the fact that our study was conducted in a setting where patients were obtaining their prescribed medications at no costs under an existing national program on universal health coverage as described elsewhere in this paper. Previous research has identified co-payments for medications as a barrier to adherence.¹³

Non-adherence to medications contributes to poor blood pressure control which subsequently increases the risk of development of end organ complications such as heart disease, chronic kidney disease, and stroke. Non-adherence to antihypertensive medications also contributes to increased healthcare costs through frequent hospitalizations related to both acute and long-term complications of uncontrolled blood pressure.²³ Therefore, it is vital that the high level of non-adherence noted in this patient population is identified as an urgent public health priority.

In this study, it was also established that nearly two-thirds of the patients on follow-up had poor hypertension knowledge, and this factor was an independent predictor of medication non-adherence. Patients with poor knowledge about hypertension were 5.6 times more likely to be non-adherent to their prescribed medications compared to those with good knowledge. Previous studies have established a similar strong link between hypertension knowledge and medication adherence.^{24,25} Findings on the responses obtained on various questions assessing specific knowledge about hypertension indicated that a significant number of patients lacked an understanding that hypertension is an asymptomatic and chronic disease requiring lifelong treatment. It is therefore likely that this knowledge deficit may have contributed to some patients skipping medications whenever they felt fine or discontinuing the medication after sometime with the assumption that they had recovered. In a survey conducted on health seeking behaviors among patients with hypertension and diabetes in diverse rural care settings in Kenya, a significant number of patients (30%) reported that they usually discontinue themselves of medication whenever they feel like they have 'recovered' from the disease.²⁶

The complexity of the dosage regimens put on patients was found to have a significant influence on medication adherence. Patients who were receiving two or more drugs were 2.8 times more likely to be non-adherent to their medications compared to those on mono-

therapy. Previous studies have also observed a similar inverse relationship between the number of prescribed medications and adherence.²⁴⁻²⁷ Having multiple medications is associated with “pill burden” and increases the likelihood of a patient forgetting to take some drugs included in his/her treatment regimen. Similarly, being on multiple daily doses (two or more doses) was also associated with a higher likelihood of non-adherence compared to receiving a single dose. An earlier meta-analysis study showed that patients on single daily doses are more likely to be adherent to their medications compared to those on higher dosing frequencies.²⁸ There is also a growing body of evidence showing that treatment regimens comprising of single pill drug combinations are associated with higher levels of adherence compared to free drug equivalent combinations.²⁹

Two important limitations need to be taken into account when making an interpretation of the findings generated from our study. Firstly, our study sample was drawn from a population comprising of patients only seen in the two selected tertiary care hospitals in Nyeri County. Given that there are other facilities in the region where hypertensive patients are followed up, our findings cannot be generalized to the entire population of hypertensive patients in Nyeri County or the country at large. However, the study provides an important entry point into quantifying the burden of medication non-adherence, and also in exploring the contributing factors to the problem among hypertensive patients on follow-up in Nyeri County. Secondly, the dependent variable of our study (medication adherence) was assessed using a self-report tool, and therefore the measurements may be viewed as an approximation on a patient's true adherence to prescribed medications. However the tool used has been validated and utilized in assessment of medication adherence in multiple previous studies in diverse settings and populations.¹⁷

Conclusion and Implications for Translation

The findings from our study indicate that medication non-adherence is highly prevalent

among hypertensive patients on follow-up in the two tertiary hospitals in Nyeri County, and therefore creating an urgent need for concerted efforts to address the problem. The factors associated with increased likelihood of non-adherence were poor knowledge about hypertension and having more complex treatment regimens. Such factors are largely modifiable and this therefore presents an opportunity for players in the healthcare delivery sector to formulate interventions to address the identified health problem. Health education based interventions can be developed and integrated in the care of hypertensive patients to enhance understanding on the various aspects of hypertension, especially the asymptomatic and chronic nature of the condition.

There is also need for policy makers to consider revising treatment guidelines to recommend where appropriate, the adoption of simpler treatment regimens comprising of single fixed drug combinations to enhance adherence.

Compliance with Ethical Standards

Conflicts of Interest:

The authors declare no competing interests.

Financial Disclosure:

Nothing to declare.

Ethics Approval:

This study was ethically reviewed and approved by Kenyatta University Ethics Review Committee [Reference Number: KU/ERC/APPROVAL/VOL. 1 (267)].

Acknowledgments:

The authors thank the medical and nursing staff of Nyeri County Referral Hospital and Consolata Mathari Mission Hospital for their invaluable support and participation during implementation of this study.

Funding/Support:

This study was funded by Dedan Kimathi University of Technology through a PhD student sponsorship grant.

References

1. **Stanaway JD, Afshin A, Gakidou E**, et al. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1923-1994.
2. **Kyu HH, Abate D, Abate KH**, et al. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1859-1922.
3. **Zhou B, Carrillo-Larco RM, Danaei G**, et al. Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet*. 2021;398(10304):957-980.
4. **Pengpid S, Peltzer K**. Prevalence, awareness, treatment and control of hypertension among adults in Kenya: cross-sectional national population-based survey. *East Mediterr Health J*. 2020;26(8):923-932.
5. **Oparil S, Acelajado MC, Bakris GL**, et al. Hypertension. *Nat Rev Dis Primers*. 2018;4:18014.
6. Kenya Ministry of Health. *Kenya National Guidelines for Cardiovascular Diseases Management*. 2018. Division of Non-Communicable Diseases. (accessed September 20, 2022)
7. World Health Organization. 2003. Adherence to Long-Term Therapies: Evidence for action. WHO; (accessed September 20, 2022)
8. **Abegaz TM, Shehab A, Gebreyohannes EA, Bhagavathula AS, Elnour AA**. Nonadherence to antihypertensive drugs: A systematic review and meta-analysis. Meta-Analysis. *Medicine*. 2017;96(4):5641.
9. **Moss JT, Kimani H, Mwanzo I**. Compliance to antihypertensive therapy and associated factors among adults' hypertensive patients attending medical clinics in Kilifi County Kenya. *Hypertension, Compliance, Antihypertensive, Therapy. IJCMPh*. 2021;8(10):11.
10. **Otenyo SO, Kereri DK**. Illness perceptions and adherence to medication regimen among hypertensive patients attending a county referral hospital in Kenya. *J Hypertens Manag*. 2021;7:59.
11. **Xiong S, Peoples N, Østbye T**, et al. Family support and medication adherence among residents with hypertension in informal settlements of Nairobi, Kenya: a mixed-method study. *J Hum Hypertens*. 2023;37(1):74-79.
12. **Achieng' L, Joshi M, Ogola E, Karari E**. Adequacy of blood pressure control and level of adherence with antihypertensive therapy. *East Afr Med J*. 2009;86(11):499-506.
13. **van der Laan DM, Elders PJM, Boons CCLM, Beckeringh JJ, Nijpels G, Hugtenburg JG**. Factors associated with antihypertensive medication non-adherence: a systematic review. *J Hum Hypertens*. 2017;31(11):687-694.

14. **Wangia E, Kandie C.** Essential elements in attaining universal health coverage in Kenya. In: Refocusing on quality of care and increasing demand for services. Government of Kenya; 2016. (accessed July 4, 2022)
15. **Nundy S, Kakar A, Bhutta ZA.** Writing the Research Proposal: The Art and the Science. *How to Practice Academic Medicine and Publish from Developing Countries?* In: A Practical Guide. Springer Singapore; 2022. p. :139-152.
16. **Sullivan KM, Dean A, Soe MM.** Open Epi: a web-based epidemiologic and statistical calculator for public health. *Public Health Rep.* 2009;124(3):471-474.
17. **Lam WY, Fresco P.** Medication Adherence Measures: An Overview. *Biomed Res Int.* 2015;2015:217047.
18. **Abu H, Aboumatar H, Carson KA, Goldberg R, Cooper LA.** Hypertension knowledge, heart healthy lifestyle practices and medication adherence among adults with hypertension. *Eur J Pers Cent Healthc.* 2018;6(1):108-114.
19. **Olowe OA, Ross AJ.** Knowledge, adherence and control among patients with hypertension attending a peri-urban primary health care clinic, KwaZulu-Natal. Observational Study. *Afr J Prim Health Care Fam Med.* 2017;9(1):e1-e7.
20. **Algabbani FM, Algabbani AM.** Treatment adherence among patients with hypertension: findings from a cross-sectional study. *Clin Hypertens.* 2020;26:18.
21. **Otenyo S, Maranga A.** Factors affecting adherence to antihypertensive medication regimen among hemodialysis patients attending a private hospital in Mombasa, Kenya. *International Journal of Pharmaceutical Sciences and Research.* 2018;12(9):755-760.
22. **Kubo MN, Kayima JK, Were AJ, Mc Ligeyo SO, Ogola EN.** Factors associated with uncontrolled hypertension among renal transplant recipients attending nephrology clinics in Nairobi, Kenya. *J Transplant.* 2015;2015:746563.
23. **Vrijens B, Antoniou S, Burnier M, de la Sierra A, Volpe M.** Current situation of medication adherence in hypertension. *Front Pharmacol.* 2017;8:100.
24. **Algabbani FM, Algabbani AM.** Treatment adherence among patients with hypertension: findings from a cross-sectional study. *Clin Hypertens.* 2020;26:18.
25. **Jankowska-Polańska B, Chudiak A, Uchmanowicz I, Dudek K, Mazur G.** Selected factors affecting adherence in the pharmacological treatment of arterial hypertension. *Patient Prefer Adherence.* 2017;11:363-371.
26. **Karinja M, Pillai G, Schlienger R, Tanner M, Ogutu B.** Care-seeking dynamics among patients with diabetes mellitus and hypertension in selected rural settings in Kenya. *Int J Environ Res Public Health.* 2019;16(11):2016.
27. **Asgedom SW, Atey TM, Desse TA.** Antihypertensive medication adherence and associated factors among adult hypertensive patients at Jimma University Specialized Hospital, southwest Ethiopia. *BMC Res Notes.* 2018;11(1):27.

28. **Coleman CI, Limone B, Sobieraj DM,** et al. Dosing frequency and medication adherence in chronic disease. *J Manag Care Pharm.* 2012;18(7):527-39.
29. **Tsioufis K, Kreutz R, Sykara G, van Vugt J, Hassan T.** Impact of single-pill combination therapy on adherence, blood pressure control, and clinical outcomes: a rapid evidence assessment of recent literature. Systematic Review. *J Hypertens.* 2020;38(6):1016-1028.